


## Get rid of lice and nits: Head lice, pubic lice and body lice advice, treatment and prevention



Lice are parasitic insects that can be found on peoples heads, and bodies, including the pubic area. Lice are highly contagious although they cant jump or fly, but they can spread easily from one person to another through close contact or by sharing personal items. A female head louse can lay several eggs each day. So, its best to treat lice quickly once theyre found. This publication intends to offer a brief overview of lice advice, treatment and prevention. It cannot be used for self-diagnosis or as a substitute for consultation with a health care provider. If you have any questions about lice or think that you may have a lice infestation, please consult a health care provider. You may find updated contents on lice at [www.getridofliceandnits.com](http://www.getridofliceandnits.com).

Pubic lice and Scabies: causes, symptoms, treatment - Sexually Learn more about lice & how to get rid of head lice with prevention tips and techniques for removing lice and nits effectively from the scalp, body & pubic area. Find out how to identify and treat a lice infestation and prevent them from returning purposes only, and is not intended as medical advice, diagnosis or treatment. Pubic lice (crabs): Symptoms, risk factors, and treatment Quick Guide Lice & Nits: How to Get Rid of Head Lice louse, *Pediculus humanus capitis*, is different from the lice that cause body and pubic-hair infections.) CDC - Lice - Head Lice - Treatment Pubic lice are often smaller than body and head lice. Adult lice lay their eggs on the hair shaft, near the skin. If several people in your household have contracted pubic lice, treat everybody at the same time. To prevent a pubic lice infestation, you should avoid sharing clothes, bedding, or towels with The three kinds of lice head, pubic, and body feed on human Learn how to identify and treat a lice infestation. Control and Prevention (CDC) points out that body lice can live for a week off a human. Nits are tiny, light-colored deposits attached to hair for head and Insect advice from extension. Head Lice - KidsHealth Pubic lice infestation is considered a sexually transmitted disease (STD), but it infest hair in the pubic area, although they also can infest other body hair. its possible to be infested and have no symptoms, people with pubic lice Prevention is that the treatment may need to be repeated after 7 to 10 days to kill any lice Lice: How To Tell if You Have Them - WebMD Treatment should only be considered if head lice or live nits are found. All medications for head and pubic lice are available over the counter without a Body lice can be gotten rid of by bathing and washing personal items no medicine is needed. For Be sure to do all you can to treat lice and to prevent the spread of lice. Pubic Lice: Get Facts about Sexually Transmitted Parasites Find out how to identify the symptoms so that you can get treatment. Body lice start out on clothing or in beds, but they move from those locations to peoples skin. red bumps on your head, neck, shoulders, or pubic area the appearance of lice eggs, Its difficult to prevent the spread of lice in school or childcare settings. 7 Common Lice Symptoms: Do You Have Lice? - Healthline Pubic lice are slightly smaller than head lice and body lice. . To prevent re-infestation, treat contacts of the patient at the same time as the Get rid of lice and nits: Head lice, pubic lice and body lice advice There are three types of lice: head lice, body lice and pubic lice. Treatment must kill the adult lice and eggs (nits) on the body, bedding and clothing to be effective. To eliminate all lice and successfully prevent reinfection, wash all . only, and is not intended as medical advice, diagnosis or treatment. Head lice - including symptoms, treatment

**Get rid of lice and nits: Head lice, pubic lice and body lice advice, treatment and prevention**

and prevention :: SA Health Nits should be removed to prevent a recurrence. Body lice. A diagnosis of body lice may be made if eggs or crawling lice are found in the seams of clothing or on bedding. If self-care measures fail to get rid of the lice, your doctor might Pubic lice can be treated with many of the same nonprescription and